



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2021

Rifabutin is inactivated by *Mycobacterium abscessus* Arr

Schäfle, Daniel ; Selchow, Petra ; Borer, Barbara ; Meuli, Michael ; Rominski, Anna ; Schulthess, Bettina ; Sander, Peter

Abstract: *Mycobacterium abscessus* exhibits arr (ADP-ribosyltransferase)-dependent rifampicin (RIF) resistance. In apparent contrast, rifabutin (RBT) has demonstrated promising activity in *M. abscessus* infection models implying that RBT might not be inactivated by Arr. RBT susceptibility testing of *M. abscessus* Δ arr revealed a strongly decreased minimal inhibitory concentration (MIC). Our findings therefore suggest that the efficacy of RBT might be enhanced by rendering RBT resilient to Arr-dependent modification or by blocking *M. abscessus* Arr activity.

DOI: <https://doi.org/10.1128/AAC.02215-20>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-196544>

Journal Article

Accepted Version

Originally published at:

Schäfle, Daniel; Selchow, Petra; Borer, Barbara; Meuli, Michael; Rominski, Anna; Schulthess, Bettina; Sander, Peter (2021). Rifabutin is inactivated by *Mycobacterium abscessus* Arr. *Antimicrobial Agents and Chemotherapy*, 65(3):e02215-20.

DOI: <https://doi.org/10.1128/AAC.02215-20>

Short-Form Paper: Rifabutin inactivation by Arr

1 Short-Form Paper

2 **Rifabutin is inactivated by *Mycobacterium abscessus* Arr**

3 Daniel Schäfle ^a, Petra Selchow ^a, Barbara Borer ^a, Michael Meuli ^a, Anna Rominski ^a, Bettina
4 Schulthess ^{a,b} and Peter Sander ^{a,b,#}

5 ^a Institute of Medical Microbiology, University of Zurich, Zurich, Switzerland; dschaeffe@imm.uzh.ch
6 (D.S.), pselchow@imm.uzh.ch (P. Se.), barbara.m.borer@gmail.com (B.B.), mmeuli@imm.uzh.ch
7 (M.M.), anna.rominski@gmail.com (A.R.), schulthe@imm.uzh.ch (B.S.)

8 ^b National Center for Mycobacteria, University of Zurich, Zurich, Switzerland

9 # Correspondence: psander@imm.uzh.ch

10 Running title: Rifabutin inactivation by Arr

11 **Key words:** *Mycobacterium*; *Mycobacterium abscessus*; Rifamycin; Rifabutin; Rifampicin; Resistance;
12 ADP-ribosyltransferase

13 **Abstract**

14 *Mycobacterium abscessus* exhibits *arr* (ADP-ribosyltransferase)-dependent rifampicin (RIF)
15 resistance. In apparent contrast, rifabutin (RBT) has demonstrated promising activity in *M.*
16 *abscessus* infection models implying that RBT might not be inactivated by Arr. RBT susceptibility
17 testing of *M. abscessus* Δarr revealed a strongly decreased minimal inhibitory concentration (MIC).
18 Our findings therefore suggest that the efficacy of RBT might be enhanced by rendering RBT
19 resilient to Arr-dependent modification or by blocking *M. abscessus* Arr activity.

20

21

Short-Form Paper: Rifabutin inactivation by Arr

22 The incidence and prevalence of pulmonary disease caused by infections with non-tuberculous
23 mycobacteria (NTM) is increasing worldwide. NTM-pulmonary disease requires a long-lasting therapy
24 of up to two years which however has a success rate of less than 50% (1, 2). *Mycobacterium*
25 *abscessus* complex (MABSC) comprised of the three subspecies *Mycobacterium abscessus* subsp.
26 *abscessus*, *Mycobacterium abscessus* subsp. *bolletii* and *Mycobacterium abscessus* subsp. *massiliense*
27 (3) is characterized by its extreme drug resistance to both broad-spectrum drugs and tuberculosis-
28 specific drugs (4, 5). Consequently, treatment options are limited and novel drugs are urgently
29 needed. Unfortunately, *M. abscessus* drug-screens revealed a low hit rate limiting the development
30 of drugs with a new mode of action (6, 7). However, repurposing of existing drugs might offer new
31 treatment options (8).

32 Rifamycins act as inhibitors of the DNA-dependent RNA polymerase (9, 10). Rifampicin (RMP),
33 although highly active against *Mycobacterium tuberculosis* (critical concentration in MGIT 1.0 mg/L)
34 (11) shows very poor growth inhibition of *M. abscessus* (MIC 128 mg/L) due to the drug-modifying
35 activity of its ADP-ribosyltransferase Arr_{MAB} (12). Arr enzymes modify the C₂₃-hydroxyl group within
36 the rifamycin core structure (13). Deletion of *arr_{Mab}* in *M. abscessus* has been shown not only to
37 decrease the MIC for RMP but also for two other rifamycins, i.e. rifaximine (RFX) and rifapentine
38 (RPT) (12).

39 Recently, the RMP analogue rifabutin (RBT) was identified in an *in vitro* *M. abscessus* drug-screen and
40 verified to be active against a variety of *M. abscessus* strains (MIC \approx 3 mg/L) (14). Subsequently, RBT
41 proved to be efficacious in *in vitro* and *in vivo* models of *M. abscessus* infection while RMP lacked
42 activity (15, 16). The antibacterial *in vitro* and *in vivo* activity suggests that RBT unlike the other
43 rifamycins might not be a substrate for Arr_{MAB} although MIC of RBT for mycobacterial pathogens
44 without *arr* [*M. tuberculosis* (critical concentrations RBT = 0.1 mg/L vs RMP 1.0 mg/L) and *M. avium*]
45 are generally lower than for RMP (11).

Short-Form Paper: Rifabutin inactivation by Arr

46 We conducted RBT susceptibility testing (**Supplemental Material**) of *M. abscessus* ATCC 19977T, its
47 isogenic Δarr deletion mutant and the complemented mutant Δarr_{MAB} pMV361_ arr_{MAB} (12) to resolve
48 this issue (**Table 1; Supplemental Table S1**). RFX and RMP served as a control. *M. abscessus* (wt)
49 showed a RBT MIC of 4 mg/L, while MICs for rifaximine (RFX; 32 mg/L) and particularly for RMP (128
50 mg/L) were considerably higher. The *M. abscessus* Δarr_{MAB} mutant exhibited increased susceptibility
51 to RMP, RFX and particularly towards RBT (**Table 1**). The \log_2 -transformed relative resistance ratios
52 (RRR) $MIC_{wt}/MIC_{\Delta arr}$ for RBT, RFX and RMP were 18, 4 and 8, respectively. The MICs of other drug
53 classes [amikacin (AMK), tetracycline (TET)] were not affected by the *arr* genotype (RRR = 0 – 1). The
54 wild-type MICs towards the rifamycins were restored in the complemented deletion mutant. The low
55 RBT MIC of the *M. abscessus* Δarr deletion mutant indicates that RBT is a substrate for Arr.
56 Furthermore, the lower RRR for RMP as compared to the RRR for RBT (8 vs. 18) suggests that
57 resistance determinants other than Arr might selectively inactivate RMP but not RBT.

58 RMP monooxygenases (Rox) are present in a wide variety of environmental bacteria and are
59 associated with decomposition of RMP (17). A corresponding rifamycin resistance mechanism has
60 also been proposed for *M. abscessus* (18). Using Rox genes of *Streptomyces venezuelae* (Sven_0481)
61 and *Nocardia farcinica* (Nfa_35080) as a query in a BlastP search putative *M. abscessus* orthologues
62 MAB_0857, MAB_3484 and MAB_1496c were identified (**Supplemental Table S2**). Of these,
63 MAB_1496c has recently been demonstrated to be a member of the *M. abscessus* resistome due to
64 its involvement in tetracycline oxygenation (19). Single and multiple unmarked deletion mutants
65 (**Figure 1; Supplemental Figure S1**) in *M. abscessus* ATCC 19977T and *M. abscessus* Δarr were
66 constructed by allelic replacement with suicide vectors containing PCR-amplified flanking regions of
67 the target genes (**Supplemental Table S3**) and apramycin-positive and *katG* (INH^S)-negative selection
68 markers (20, 21) (**Fig. 1**). Mutants were confirmed by PCR and Southern blot analysis (**Supplemental**
69 **Figure S1**) and tested for rifamycin, tetracycline and amikacin susceptibility (**Table 1**). Exploratory
70 investigations indicated that neither individual deletion of MAB_0857, MAB_3483 and MAB_1496c
71 nor combined deletion of MAB_0857 and MAB_3483 in *M. abscessus* ATCC 19977T affected the RBT,

Short-Form Paper: Rifabutin inactivation by Arr

72 RFX, RMP and AMK MIC, respectively. Likewise, deletion of these genes in *M. abscessus* Δarr_{MAB} did
73 not alter susceptibility towards these antibiotics as compared to *M. abscessus* Δarr_{MAB} indicating that
74 these genes do not affect rifamycins. Deletion of MAB_1496c, encoding monooxygenase TetX
75 decreased the tetracycline MIC 16 -32-fold, while rifamycin MICs were not affected (for the MIC of
76 the most advanced strains see **Table 1**).

77 So far, our investigations have not identified genetic determinants responsible for selective RMP
78 modification. A phenotypic screen of a genome wide transposon mutant library might be the most
79 efficient way to identify those resistance determinants (22, 23). However, our findings on RBT
80 susceptibility of the Δarr mutant are of major importance for future drug development. The RRR of
81 about 250.000 ($\log_2=18$) clearly shows that RBT is a substrate of Arr which in turn suggests that the
82 potency of RBT against *M. abscessus* might be improved by either Arr-inhibitors or by RBT
83 modification. Regimens co-applying drugs in conjunction with inhibitors of drug-modifying enzymes
84 have been developed to restore antibiotic activity and are widely used in clinics (24-26). Similarly,
85 Arr-inhibitors might be beneficial to establish a rifamycin-based treatment for *M. abscessus*
86 infections. Alternatively modifications of the drug itself might render RBT resilient towards Arr. E.g.
87 substitution of the acetyl moiety by a bulky residue at position C₂₅ rendered RMP partially resistant
88 towards Arr-modification (12, 27, 28). We envision that these approaches may improve treatment
89 options and outcomes of pulmonary disease caused by the opportunistic pathogen feared for its
90 high-level drug resistance, *M. abscessus*.

91

92 **Authors contributions**

93 Conceptualization: P.Sa., B.S., A.R.; investigations: D.S., P.Se, B.B., M.M., A.R.; writing – original draft
94 preparation D.S., P.Sa; writing – review and editing, P.Sa. All authors have read and agreed to the
95 published version of the article.

96

Short-Form Paper: Rifabutin inactivation by Arr

97 **Funding**

98 This research was funded by Swiss National Science Foundation 310030_197699), Cystic Fibrosis
99 Switzerland (CFCH), Stiftung wissenschaftliche Forschung (University of Zurich; STWF-18-011), Swiss
100 Lung Association and Georg and Bertha Schwyzer-Winiker Stiftung (2018-02-Sa).

101

102 **Acknowledgements**

103 We would like to acknowledge generous support from the Institute of Medical Microbiology and
104 University of Zurich.

105

106 **Conflict of interest**

107 The authors declare no conflict of interest.

108

109

110 **References**

- 111 1. Griffith DE, Aksamit T, Brown-Elliott BA, Catanzaro A, Daley C, Gordin F, Holland SM,
112 Horsburgh R, Huitt G, Iademarco MF, Iseman M, Olivier K, Ruoss S, Reyn CFv, Richard J.
113 Wallace J, Winthrop K. 2007. An Official ATS/IDSA Statement: Diagnosis, Treatment, and
114 Prevention of Nontuberculous Mycobacterial Diseases. *American Journal of Respiratory and*
115 *Critical Care Medicine* 175:367-416.
- 116 2. Jarand J, Levin A, Zhang L, Huitt G, Mitchell JD, Daley CL. 2011. Clinical and Microbiologic
117 Outcomes in Patients Receiving Treatment for Mycobacterium abscessus Pulmonary Disease.
118 *Clinical Infectious Diseases* 52:565-571.
- 119 3. Tortoli E, Kohl TA, Brown-Elliott BA, Trovato A, Leao SC, Garcia MJ, Vasireddy S, Turenne CY,
120 Griffith DE, Philley JV, Baldan R, Campana S, Cariani L, Colombo C, Taccetti G, Teri A, Niemann
121 S, Wallace RJ, Jr., Cirillo DM. 2016. Emended description of Mycobacterium abscessus,
122 Mycobacterium abscessus subsp. abscessus and Mycobacterium abscessus subsp. bolletii and
123 designation of Mycobacterium abscessus subsp. massiliense comb. nov. *Int J Syst Evol*
124 *Microbiol* 66:4471-4479.
- 125 4. Luthra S, Rominski A, Sander P. 2018. The Role of Antibiotic-Target-Modifying and Antibiotic-
126 Modifying Enzymes in Mycobacterium abscessus Drug Resistance. *Frontiers in Microbiology*
127 9.
- 128 5. Johansen MD, Herrmann JL, Kremer L. 2020. Non-tuberculous mycobacteria and the rise of
129 Mycobacterium abscessus. *Nature Reviews Microbiology* 18:392-407.
- 130 6. Dal Molin M, Selchow P, Schafle D, Tschumi A, Ryckmans T, Laage-Witt S, Sander P. 2019.
131 Identification of novel scaffolds targeting Mycobacterium tuberculosis. *J Mol Med (Berl)*
132 97:1601-1613.
- 133 7. Malin JJ, Winter S, van Gumpel E, Plum G, Rybniker J. 2019. Extremely Low Hit Rate in a
134 Diverse Chemical Drug Screen Targeting Mycobacterium abscessus. *Antimicrobial Agents and*
135 *Chemotherapy* 63.
- 136 8. Wu ML, Aziz DB, Dartois V, Dick T. 2018. NTM drug discovery: status, gaps and the way
137 forward. *Drug Discovery Today* 23:1502-1519.
- 138 9. Wehrli W, Knusel F, Staeheli M. 1968. Action of Rifamycin on Rna-Polymerase from Sensitive
139 and Resistant Bacteria. *Biochemical and Biophysical Research Communications* 32:284-&.
- 140 10. Campbell EA, Korzhova N, Mustaev A, Murakami K, Nair S, Goldfarb A, Darst SA. 2001.
141 Structural mechanism for rifampicin inhibition of bacterial RNA polymerase. *Cell* 104:901-
142 912.
- 143 11. Cambau E, Viveiros M, Machado D, Raskine L, Ritter C, Tortoli E, Matthys V, Hoffner S, Richter
144 E, Del Molino MLP, Cirillo DM, van Soolingen D, Bottger EC, Grp EES. 2015. Revisiting
145 susceptibility testing in MDR-TB by a standardized quantitative phenotypic assessment in a
146 European multicentre study. *Journal of Antimicrobial Chemotherapy* 70:686-696.
- 147 12. Rominski A, Roditscheff A, Selchow P, Bottger EC, Sander P. 2017. Intrinsic rifamycin
148 resistance of Mycobacterium abscessus is mediated by ADP-ribosyltransferase MAB_0591.
149 *Journal of Antimicrobial Chemotherapy* 72:376-384.
- 150 13. Baysarowich J, Koteva K, Hughes DW, Ejim L, Griffiths E, Zhang K, Junop M, Wright GD. 2008.
151 Rifamycin antibiotic resistance by ADP-ribosylation: Structure and diversity of Arr. *Proc Natl*
152 *Acad Sci U S A* 105:4886-91.
- 153 14. Aziz DB, Low JL, Wu ML, Gengenbacher M, Teo JWP, Dartois V, Dick T. 2017. Rifabutin Is
154 Active against Mycobacterium abscessus Complex. *Antimicrobial Agents and Chemotherapy*
155 61.
- 156 15. Dick T, Shin SJ, Koh WJ, Dartois V, Gengenbacher M. 2020. Rifabutin Is Active against
157 Mycobacterium abscessus in Mice. *Antimicrob Agents Chemother* 64.
- 158 16. Johansen MD, Daher W, Roquet-Baneres F, Raynaud C, Alcaraz M, Maurer FP, Kremer L.
159 2020. Rifabutin Is Bactericidal against Intracellular and Extracellular Forms of Mycobacterium
160 abscessus. *Antimicrobial Agents and Chemotherapy* 64.

Short-Form Paper: Rifabutin inactivation by Arr

- 161 17. Koteva K, Cox G, Kelso JK, Surette MD, Zubyk HL, Ejim L, Stogios P, Savchenko A, Sorensen D,
162 Wright GD. 2018. Rox, a Rifamycin Resistance Enzyme with an Unprecedented Mechanism of
163 Action. *Cell Chemical Biology* 25:403-+.
- 164 18. Ganapathy US, Dartois V, Dick T. 2019. Repositioning rifamycins for Mycobacterium
165 abscessus lung disease. *Expert Opinion on Drug Discovery* 14:867-878.
- 166 19. Rudra P, Hurst-Hess K, Lappierre P, Ghosh P. 2018. High Levels of Intrinsic Tetracycline
167 Resistance in Mycobacterium abscessus Are Conferred by a Tetracycline-Modifying
168 Monooxygenase. *Antimicrobial Agents and Chemotherapy* 62.
- 169 20. Dal Molin M, Gut M, Rominski A, Haldimann K, Becker K, Sander P. 2018. Molecular
170 Mechanisms of Intrinsic Streptomycin Resistance in Mycobacterium abscessus. *Antimicrobial*
171 *Agents and Chemotherapy* 62.
- 172 21. Rominski A, Selchow P, Becker K, Brulle JK, Dal Molin M, Sander P. 2017. Elucidation of
173 Mycobacterium abscessus aminoglycoside and capreomycin resistance by targeted deletion
174 of three putative resistance genes. *Journal of Antimicrobial Chemotherapy* 72:2191-2200.
- 175 22. Foreman M, Gershoni M, Barkan D. 2020. A Simplified and Efficient Method for Himar-1
176 Transposon Sequencing in Bacteria, Demonstrated by Creation and Analysis of a Saturated
177 Transposon-Mutant Library in Mycobacterium abscessus. *mSystems* 5.
- 178 23. Laencina L, Dubois V, Le Moigne V, Viljoen A, Majlessi L, Pritchard J, Bernut A, Piel L, Roux AL,
179 Gaillard JL, Lombard B, Loew D, Rubin EJ, Brosch R, Kremer L, Herrmann JL, Girard-Misguich
180 F. 2018. Identification of genes required for Mycobacterium abscessus growth in vivo with a
181 prominent role of the ESX-4 locus. *Proc Natl Acad Sci U S A* 115:E1002-E1011.
- 182 24. Wise R, Andrews JM, Bedford KA. 1978. Invitro Study of Clavulanic Acid in Combination with
183 Penicillin, Amoxycillin, and Carbenicillin. *Antimicrobial Agents and Chemotherapy* 13:389-
184 393.
- 185 25. Drawz SM, Papp-Wallace KM, Bonomo RA. 2014. New beta-lactamase inhibitors: a
186 therapeutic renaissance in an MDR world. *Antimicrob Agents Chemother* 58:1835-46.
- 187 26. Kaushik A, Ammerman NC, Parrish NM, Nueremberger EL. 2019. New beta-Lactamase
188 Inhibitors Nacubactam and Zidebactam Improve the In Vitro Activity of beta-Lactam
189 Antibiotics against Mycobacterium abscessus Complex Clinical Isolates. *Antimicrob Agents*
190 *Chemother* 63.
- 191 27. Combrink KD, Denton DA, Harran S, Ma Z, Chapo K, Yan D, Bonventre E, Roche ED, Doyle TB,
192 Robertson GT, Lynch AS. 2007. New C25 carbamate rifamycin derivatives are resistant to
193 inactivation by ADP-ribosyl transferases. *Bioorganic & Medicinal Chemistry Letters* 17:522-
194 526.
- 195 28. Combrink KD, Ramos AR, Spring S, Schmidl S, Elizondo K, Morin P, De Jesus B, Maurer FP.
196 Rifamycin derivatives active against pathogenic rapidly-growing mycobacteria.
197 *Bioorganic & Medicinal Chemistry Letters* 29:2112-2115.

198

199

200 **Table 1: Minimal Inhibitory Concentration (MIC)* and Relative Resistance Ratios (RRR)** of strains**

Strain	RBT		RFX		RMP		TET		AMK	
	MIC	RRR	MIC	RRR	MIC	RRR	MIC	RRR	MIC	RRR
<i>M. abscessus</i>	4	/	32	/	128	/	32	/	1	/
Δarr_{MAB}	1.1×10^{-5}	18	1.5	4	0.38	8	24	0	1	0
Δarr_{MAB} <i>pMV361-arr_{Mab}</i>	4	0	16	1	64	1	16	1	1	0
Δarr_{MAB} <i>ΔMAB_{0857}</i> <i>ΔMAB_{3483}</i>	4.6×10^{-5}	16	1.5	4	0.25	9	24	0	1	0
Δarr_{MAB} <i>ΔMAB_{1496c}</i>	4.6×10^{-5}	16	1.5	4	0.19	9	1	5	1	0


201 *MIC (median of 6 replicates) in mg/L; **Relative Resistance Ratio (RRR) \log_2 MIC_{wt}/MIC_{comparator}

202

203

204

Short-Form Paper: Rifabutin inactivation by Arr



205

206 **Figure 1**

207 Genealogy of strains

208